

MEDICATION INFORMATION LEAFLET FOR

ETOPOSIDE

FOR INTRAVENOUS ADMINISTRATION



This leaflet should be used in conjunction with the complete consumer medicine information (CMI) which is available from any pharmacy.

This leaflet contains information about Etoposide and Etopophos® including details of known side effects. It does not include every possible side effect.

The type and severity of side effects experienced differs for each child. Some side effects are minor and easily treated at home. However, others may be more serious and require prompt medical assessment. Side effects from treatment are routinely monitored with blood tests and clinical examination. If you have any questions about side effects during treatment, please speak to your doctor. Also, if your child experiences any problems or side effects during treatment, please contact your doctor, clinical nurse consultant/coordinator or pharmacy staff for advice.

TWO TYPES

- Etoposide
- Etopophos® (etoposide phosphate)

OTHER NAMES

- VP-16

PRESENTATION AND ADMINISTRATION

- Etoposide is a clear, slightly yellow liquid which is diluted in a bag of fluid for intravenous infusion.
- Etopophos® (etoposide phosphate) is the water soluble form of etoposide. It is a clear, colourless liquid which is either drawn up in a syringe or diluted in a bag of fluid. It can be given by intravenous injection or intravenous infusion.

SIDE EFFECTS

Common

IMMEDIATE (within 1-2 days of receiving drug)	DELAYED (within 1-2 weeks of receiving drug)
<ul style="list-style-type: none">• Nausea• Vomiting	<ul style="list-style-type: none">• Decrease in blood count• Hair loss

Less common

- Low blood pressure
- Loss of appetite
- Mucositis (inflammation or ulceration of the mouth and gut)
- Diarrhoea

Rare

- Allergic reactions which may be life-threatening. Symptoms of a severe reaction may include skin rash or hives, swelling or tingling of the lips and face, difficulty breathing and low blood pressure which may lead to collapse. Seek medical attention **immediately** if any of the above symptoms occur.
- Numbness and tingling in hands and feet.
- Skin rash.
- New leukaemia or tumour (very rare)

ADDITIONAL INFORMATION

- Avoid excessive exposure to sunlight and use a high SPF sunscreen.
- Keep ALL medication out of the reach of children.

CONTACT

For further information, or to discuss any concerns you may have about your child's medication, please feel free to contact the Children's Cancer Centre of your child's treating hospital:

- The Royal Children's Hospital Oncology Pharmacy (03) 9345 6290
- Monash Children's Hospital Cancer Centre (03) 8572 3456 or Pharmacist desk (03) 8572 3464

Disclaimer: This information has been prepared by the Paediatric Integrated Cancer Service, with input from the Children's Cancer Centre Pharmacy Departments at The Royal Children's Hospital and Monash Children's Hospital. This information is for educational purposes only and should not be seen as a substitute for advice from your doctor or other professional healthcare provider. If you have specific questions about the content of this information, or any other medical matter, it is recommended that you consult your doctor, pharmacist or other professional healthcare provider. This information is considered to be true and correct at the date of publication, however changes in circumstances after the time of publication may impact on the accuracy of this information.

Reviewed March 2017

Available at www.pics.org.au/families